

TOPICAL REVIEW • **OPEN ACCESS**

# A network-based approach for semi-quantitative knowledge mining and its application to yield variability

To cite this article: Bernhard Schauburger *et al* 2016 *Environ. Res. Lett.* **11** 123001

View the [article online](#) for updates and enhancements.

## Related content

- [Wheat yield loss attributable to heat waves, drought and water excess at the global, national and subnational scales](#)  
M Zampieri, A Ceglar, F Dentener et al.
- [Heat stress is overestimated in climate impact studies for irrigated agriculture](#)  
Stefan Siebert, Heidi Webber, Gang Zhao et al.
- [The impact of climate extremes and irrigation on US crop yields](#)  
T J Troy, C Kipgen and I Pal

## Environmental Research Letters



## TOPICAL REVIEW

## OPEN ACCESS

## RECEIVED

7 October 2016

## ACCEPTED FOR PUBLICATION

11 November 2016

## PUBLISHED

28 November 2016

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



# A network-based approach for semi-quantitative knowledge mining and its application to yield variability

**Bernhard Schaubberger<sup>1</sup>, Susanne Rolinski and Christoph Müller**

Potsdam Institute for Climate Impact Research, Telegraphenberg A56, D-14473 Potsdam, Germany

<sup>1</sup> Author to whom any correspondence should be addressed.E-mail: [schauber@pik-potsdam.de](mailto:schauber@pik-potsdam.de), [rolinski@pik-potsdam.de](mailto:rolinski@pik-potsdam.de) and [cmueller@pik-potsdam.de](mailto:cmueller@pik-potsdam.de)**Keywords:** yield variability, crop models, interaction network, plant process, wheat, maize, riceSupplementary material for this article is available [online](#)


## Abstract

Variability of crop yields is detrimental for food security. Under climate change its amplitude is likely to increase, thus it is essential to understand the underlying causes and mechanisms. Crop models are the primary tool to project future changes in crop yields under climate change. A systematic overview of drivers and mechanisms of crop yield variability (YV) can thus inform crop model development and facilitate improved understanding of climate change impacts on crop yields. Yet there is a vast body of literature on crop physiology and YV, which makes a prioritization of mechanisms for implementation in models challenging. Therefore this paper takes on a novel approach to systematically mine and organize existing knowledge from the literature. The aim is to identify important mechanisms lacking in models, which can help to set priorities in model improvement. We structure knowledge from the literature in a semi-quantitative network. This network consists of complex interactions between growing conditions, plant physiology and crop yield. We utilize the resulting network structure to assign relative importance to causes of YV and related plant physiological processes. As expected, our findings confirm existing knowledge, in particular on the dominant role of temperature and precipitation, but also highlight other important drivers of YV. More importantly, our method allows for identifying the relevant physiological processes that transmit variability in growing conditions to variability in yield. We can identify explicit targets for the improvement of crop models. The network can additionally guide model development by outlining complex interactions between processes and by easily retrieving quantitative information for each of the 350 interactions. We show the validity of our network method as a structured, consistent and scalable dictionary of literature. The method can easily be applied to many other research fields.

## 1. Introduction

Crop yields can vary strongly between years and locations. These fluctuations, or yield variability (YV), are undesirable, since they undermine food security on three dimensions (Morton 2007, Schmidhuber and Tubiello 2007, Wheeler and von Braun 2013, Thornton *et al* 2014). First, the amount of harvested food can be lower than necessary, second, the financial sustainability of farming systems can be challenged, and third, the access to nutritious food can be diminished by rising prices or export bans connected to variable yields (Headey and Fan 2008, Headey 2010, Coumou and Rahmstorf 2012, Chung *et al* 2014). Substantial

fractions of historic YV can be explained by weather variability and extremes like droughts, floods, heat waves, cold spells, or combinations of them (Porter and Semenov 2005, Schlenker and Roberts 2009, Coumou and Rahmstorf 2012, Lobell *et al* 2013, Der-yng *et al* 2014, Ray *et al* 2015, Lesk *et al* 2016). Globally about one third of YV can be explained by weather variation, but with large regional differences (Ray *et al* 2015). Although some of the actual weather-induced variation in yields might be lost in the aggregation procedure, this leaves up to two thirds of YV to be explained (SI figure S1). Thus other environmental or management factors must cause the variation. An example for regional differences is the



influence of precipitation on YV (figure 1). Precipitation variability clearly determines wheat variability in Australia (panel (a) with inset), while in Germany wheat does not exhibit a clear, simple response to either temperature or precipitation (panel (b); Ray *et al* 2015).

Variability in growing conditions is transmitted to yield levels by plant physiological processes. These form a layer of complexity that has to be accounted for when assessing future YV. A huge body of experimentally-derived knowledge describes quantitative relationships between growing conditions, plant physiological processes and yield (e.g. Barnabás *et al* 2008, Farooq *et al* 2009b, Hatfield *et al* 2011). Process-based crop models are frequently used to study the influence of growing conditions on crop physiology and yields apart from experiments. These models represent our current knowledge on plant interactions with their environment (Boote *et al* 2013, Holzworth *et al* 2015). They are apt to reliably reproduce spatially aggregated mean yield levels (Palosuo *et al* 2011, Asseng *et al* 2015, Martre *et al* 2015).

Despite the abundant knowledge about YV a consistent and comprehensive overview of its causes and mechanisms is not yet available. Apart from the study by Ray *et al* (2015) and similar predecessors (see references therein) other causes of YV were also researched, but focusing on subsets of possible causes only. Bakker *et al* (2005) decipher the contribution of soil, climate and management as important sources of spatial wheat YV in Europe. Porter and Semenov (2005) or Asseng *et al* (2011) consider the impacts of heat stress on crop yields, but do not consider other climatic

factors like water or solar radiation, or do not discuss plant physiological processes. Other studies include Yu *et al* (2014), who identify temporal patterns of climate effects on wheat YV in Australia but do not consider processes, or Thornton *et al* (2014), who stress the importance of considering climate variability in food security assessments, and Ben-Ari and Makowski (2014), who identify the geographical distribution of crops as source of YV. At the same time crop models are deemed to lack adequate implementations of temporal YV under changing growing conditions (Rötter *et al* 2011, Sánchez *et al* 2014). In particular, extreme events like heat or drought have been found to be less well represented (Palosuo *et al* 2011, Rötter *et al* 2011, White *et al* 2011, Boote *et al* 2013, Rötter 2014, Asseng *et al* 2015). A comprehensive overview of the current status of crop models is provided by Boote *et al* (2013), who list nine cardinal points on how to improve crop models. Yet YV is not explicitly addressed as a topic, and stresses are only considered for heat, nitrogen and water. In Holzworth *et al* (2015) the authors state the effects of increased CO<sub>2</sub>, temperature extremes, pests and hydrology as inadequately represented in models. Barlow *et al* (2015) and Eyshi Rezaei *et al* (2014) describe the negative effects of frost or heat on cereals and derive modeling guidelines. We conclude that a comprehensive and systematic overview of causes and mechanisms of YV is much needed, in particular for selecting suitable process candidates for model improvement.

Therefore we systematically review the literature on YV, and provide specific recommendations on how to incorporate the findings into process-based crop

models. We adopt a novel, semi-quantitative technique for systematic reviews since the literature on plant physiology is overwhelming (more than 11 000 hits in the

absolute yield amounts (e.g. t/ha) between growing seasons of the same crop at the same location; an example measure would be the standard deviation.

$$\Delta\text{Yield} := \Delta \left[ \int_{\text{growing season}} \text{physiological processes (growing conditions)} \right]. \quad (2.1)$$

Web of Science<sup>2</sup> database for ‘crop variability’). The idea is to structure knowledge in a network of interactions, where management, weather and other environmental factors define crop yield via plant physiological processes (figure 3). We then rank the possible contribution of individual growing parameters to yield from their location in the network topology, independent from their frequency in the literature. Furthermore, we quantify the importance of plant physiological processes for the transmission of variability in growing conditions to YV. Finally, we use this knowledge to compile suggestions for the improved representation of YV in crop models. The method is ‘semi-quantitative’ since we do not employ quantitative relationships between growing conditions and yield. But we do quantify the impact of growing condition parameters and plant physiological processes by their contribution to the network structure. To test the validity of the method we compare our proposals to the agenda suggested by Boote *et al* (2013). We consider maize (a  $C_4$  plant), rice and wheat (both  $C_3$  plants), representing roughly 92% of the globally harvested cereals (Ben-Ari and Makowski 2014) and planted on 41.3% of the global agricultural area (Portmann *et al* 2010).

This article describes a new method for mining knowledge from the literature, which is applied to review physiological mechanisms of YV. It is bound to reproduce existing knowledge to a large extent, but will check this for comprehensiveness and can thus guide future crop model development. With this review, we aim to answer three questions. First, what are key drivers of YV in wheat, maize and rice? Second, what are the central plant physiological processes involved? Third, how can the important interactions be included into crop models?

## 2. Materials and methods

### 2.1. Definitions and network terminology

Yield is an aggregate measure of crop characteristics and performance over the entire growing season. Yield can be defined as the integral of many short-term variations in growing conditions during the growing season and the plant’s reaction towards them. Temporal YV is hence the variability of this integral (equation (2.1)). We define YV as average changes in

We focus on the fine-grain interactions between growing conditions, plant physiological processes and yield. Spatial variability plays, next to variability over time, a decisive role (Ben-Ari and Makowski 2014). Here we assume that spatial and temporal variation share common causes like e.g. temperature variation over space or time (Blois *et al* 2013), such that our analysis is also valid for spatial YV. We do not consider long-term trends, including a gradual increase in yields through improved management or a shift in yield trends from changes in climatic conditions. We use the term ‘stress’ to describe any non-optimal growing condition (e.g. a heat wave). Finally, plant growth and plant development (‘phenology’) are two distinct terms: while the first is a physical accumulation of biomass over the growing season, the latter refers to advances in developmental stages, for example the transition from vegetative to reproductive growth.

A network consists of *nodes* (i.e. elements) and *edges* between these. In our case nodes refer to processes, drivers or variables and edges to interactions between them. The *source/target node* of an edge is its starting/end point, respectively. A *path*  $q$  from node A to node B, denoted as  $A \xrightarrow{q} B$ , through the network follows a direction and can be direct (i.e. connecting A with B immediately) or indirect (i.e. containing intermediate nodes). The *path length*  $|q|$  is the number of edges it contains (illustration in SI figure S2).

### 2.2. Network construction

The starting point for the network construction was the basic network scheme shown in figure 3, into which subcategories and interactions were subsequently added. The interactions described in six standard physiology text books (Hay and Walker 1989, Porter and Lawlor 1991, Hall *et al* 1993, Larcher 1995, Hay and Porter 2006, Lambers *et al* 2008) were used to add details: the network was refined with every encountered subcategory or interaction. For example, if the scanned literature stated an influence of temperature (T) on photosynthesis, these two nodes were created (if not yet existent) and an interaction arrow drawn from T to photosynthesis (if not yet existent). If an interaction edge was already present, the new reference was recorded, but no duplicate edge was included. This ensures that interactions do not gain more weight just because they are frequently stated in the literature. With more details, the categories were

<sup>2</sup> <http://apps.webofknowledge.com/>; accessed on 11 October 2015.

subdivided. For each node and interaction it was annotated for which crop (wheat/maize/rice or all three) it is valid, thus creating crop-specific networks.

Afterwards a systematic search for studies in the full ISI Web of Science database<sup>3</sup> was performed. A keyword list with 55 entries was created, using terms from the initial textbook-based network. General terms like ‘*yield variability*’ and more specific ones like ‘*temperature AND wheat AND yield*’ were included (full keyword list in SI table S3). Only papers after 1990 were considered to limit the number of search results. This first search for the keywords in the ‘Topic’ fields yielded 460 765 studies in total, so the results were filtered to contain only ‘Review’ papers. If this number was still large ( $> 200$ ) for one search term the results were further filtered to contain the keywords in the ‘Title’ instead of ‘Topic’ (with few exceptions; SI table S3). Additionally, references to and in four large reviews (Barnabás *et al* 2008, Farooq *et al* 2009b, Hatfield *et al* 2011, Boote *et al* 2013) were searched to validate the efficacy of the keyword approach. These search criteria resulted in 8818 studies that were inspected for relevance by sequentially looking at title, abstract and full text. An article was *relevant* if the study included an explicit treatment of plant physiological processes, with either growing condition influences on them or their influence on yield, and the interactions were not derived solely from modeling studies. More recent studies were selected when similar but older ones existed. Molecular details like enzyme activity or signaling molecules and genotypic or cultivar-specific differences are not considered. After this final filter step, 60 relevant papers remained from which interactions were manually included in the initial text book-based network.

Six out of 350 edges were added without explicit literature reference as they were considered obvious but have not been found in the selected literature. These are: irrigation adds to soil water content (SWC), fertilization adds to soil nutrient levels, sowing and harvesting time affect the amount of precipitation and solar radiation intercepted during the growing season, water uptake is affected by SWC, and the plant’s uptake of micronutrients influences their content in grains.

### 2.3. Driver and process importance

The importance of drivers as possible sources of YV was derived from the network structure. The importance of a driver  $d$  is defined by the number of different paths from  $d$  to yield amount, mediated by various plant physiological processes (equation (2.2);  $m$  denotes the maximum path length).

$$\text{importance}_m(d) := |\{q \mid d \xrightarrow{q} \text{Yield} \wedge |q| \leq m\}|. \quad (2.2)$$

A maximum path length  $m = 4$  (i.e. at most three intermediate nodes) was chosen. This allows for possibly important indirect effects but avoids cyclic paths. Sensitivity to this assumption was tested with path lengths from 1 to 10. Each interaction was counted only once, independent of the number of studies which mentioned that specific influence. Thus a frequent occurrence of an interaction in the literature does not necessarily imply a high ranking. It is assumed that only drivers that exert substantial impact on plant physiology *and* are variable in nature can cause YV. Therefore each possible driver was qualitatively classified for its variability in nature and drivers with low variation were excluded. Three reduced network variants were also analyzed to search for variability drivers other than temperature (T) and precipitation (Pr). From the full network either T (air and soil, with all out-edges), or Pr along with SWC, or both T and Pr nodes (then also air humidity) were deleted; then the importance assessment was repeated.

The importance of a process for transmitting variability in growing conditions to YV was evaluated by plotting both impact values against each other (scheme in figure 2). The *impact* of a node  $v$  on another node  $w$  is defined by the number of paths between  $v$  and  $w$ , similar to the importance of drivers in equation (2.2). A process which is impacted by many different influences from the growing conditions (above the mean value on  $x$ -axis) and in turn substantially impacts yield levels (above mean value of  $y$ -axis) was assumed *important* in this respect. Processes in the other sectors of the plot did fulfill either one criterion or none at all, and were thus deemed less or not important for shaping yield amounts.

We consider this network method as ‘semi-quantitative’ since quantitative relationships between growing conditions and yield are not included, but the relative impact of growing condition parameters and plant physiological processes is evaluated by their quantitative contribution to the network structure.



The network approach explicitly integrates across physiological scales and assumes that driver or process importance is directly related to their number of network links to yield. The adequacy of these two assumptions is justified in the discussion section.

## 3. Results

### 3.1. Network structure

The crop yield interaction network contains 130 nodes and 509 edges. Of the edges 350 are interactions between nodes (*functional interactions*); the other edges only connect hierarchical distinctions in categories, e.g. ‘uptake’ to ‘uptake of nutrients’. Each node is connected on average by 3.92 edges (functional edges only: 2.69), the average number of studies cited

<sup>3</sup> <http://apps.webofknowledge.com/>; accession dates in SI table S3.




per functional interaction is 1.53, and the nodes with the highest out-degree are temperature (49 out-edges), SWC (39) and CO<sub>2</sub> (26). The number of edge annotations for only wheat are 105, for only maize 36 and for only rice 32; interaction references applying to all three crops summarize to 363 (SI table S1). A condensed version of the network is shown in figure 3 where interaction and citation numbers are split between categories. The full interaction network is provided in the SI (figure S3 and as GraphML editable network file). Among the drivers of YV in growing conditions we considered the following stressors:

chilliness and heat, water logging and drought, excess and shortage of solar radiation, ozone, strong wind, nutrient shortage and excess, salt and acidity stress, pests and diseases, and toxic substances.

### 3.2. Relative importance of factors causing YV

SWC, with its climatic precursor precipitation (Pr), and temperature (T) are ranked as foremost influences on yield by our method. An analysis of the full and the reduced network variants suggests also the following environmental factors as physiologically important for





yield amount (but not necessarily its variability): carbon dioxide, solar radiation, soil salinity, tropospheric ozone concentration, microorganisms (e.g. mycorrhizas), soil temperature, soil pH, soil density, wind and soil nutrient levels. From the management category the following nodes are suggested as important: timing of sowing/harvesting, weed management, irrigation, soil management and drought resistance support. Figure 4 shows the relative ranking of drivers (only top 25%): the  $x$ -axis contains the four network types (full and three reduced variants) and the  $y$ -axis the number of interactions up to a path length of four (log-scale). The more interactions a factor controls, the more important it is assumed for yield formation. The results are similar for all three crops, although the relative importance can be crop-specific (SI figure S7). Different thresholds for maximum path lengths do not change the results qualitatively (SI figure S4).

### 3.2.1. Filtering drivers with low short-term variability

Only factors that are variable in nature can be drivers of variable yields. Therefore, to exclude unlikely drivers of variability, we determine the variation of each factor that is regarded as yield-influencing from our network. Table 1 lists the variability of each factor and whether it is considered in this review. The management options listed above are 'variable' by definition since the farmer can decide at any point in time to apply irrigation, drought support (seed priming only before the growing season), weed control or different soil management options (before and within season). Sowing times can also be highly variable between years, depending on local climatic conditions, cultivar choice, soil parameters and other factors

(Craufurd and Wheeler 2009, Portmann *et al* 2010, Waha *et al* 2012, 2013). The impact of management decisions on YV is not assessed here, but should nonetheless be considered in crop models. In the following we only consider environmental variations as source of YV. Interactions between drivers and plant physiological processes are summarized in cursory depth in the next section. An extended and in-depth version with more references can be found in the SI.

### 3.2.2. Processes affected by water and temperature

The influence of precipitation on yield is paramount in most regions of the globe (Yu *et al* 2014, Ray *et al* 2015), and it is mediated via the SWC. SWC depends on precipitation and other factors like temperature, soil density and management (e.g. tillage) (Leakey *et al* 2009, Hatfield *et al* 2011). The fraction of SWC that is available for uptake by plant roots is further determined by soil salinity or competition (Fuhrer 2003, Tokatlidis 2014). Photosynthesis, temperature regulation, carbon allocation, nutrient uptake and reproduction strongly depend on water to function properly (Boyer and Westgate 2004, Reddy *et al* 2004, Barnabás *et al* 2008, Brouder and Vole nec 2008, Farooq *et al* 2009b, Gonzalez-Dugo *et al* 2010, Ahmed *et al* 2013, Jagadish *et al* 2014, Suzuki *et al* 2014). In particular, reproductive processes including anthesis and grain filling are highly sensitive to drought (Acevedo *et al* 2002, Boyer and Westgate 2004, Barnabás *et al* 2008, Lawlor and Tezara 2009, Gonzalez-Dugo *et al* 2010, Thitisaksakul *et al* 2012, Powell *et al* 2012, Ashraf 2014, Farooq *et al* 2014, Jagadish *et al* 2014). Non-optimal water availability

**Table 1.** Assessment of the natural variability of important yield-influencing factors in crop growing conditions. The first column contains the factor, the second column its short-term variability in nature (low or high), the third column lists references for the variability, the fourth column contains comments on the factor and the fifth states if the factor is included in this review.

Factor	Variability	References	Comment	Inclusion
Soil water content and precipitation	High	Lobell and Gourdji (2012), Donat <i>et al</i> (2013), Ruane <i>et al</i> (2015)	SWC buffers Pr variability, but eventually follows the Pr trend (Bell <i>et al</i> 2010)	Yes
Temperature (air and soil)	High	Rahmstorf and Coumou (2011), Seneviratne <i>et al</i> (2012)	For example influences on yield see Ray <i>et al</i> (2015)	Yes
Solar radiation	High	Wang and Dickinson (2013)	Important especially when other factors are not limiting (Tollenaar and Lee 2002, de Bossor-eille de Ribou <i>et al</i> 2013)	Yes
Tropospheric Ozone	High	Fuhrer (2003), Martiello and Giacchi (2010), Wild <i>et al</i> (2012), Tai <i>et al</i> (2014), Hoshika <i>et al</i> (2015)	Ozone follows temperature, solar radiation and precursor trends nonlinearly McGrath <i>et al</i> (2015)	Yes
Wind	High	SI figure S3 for interactions	Aggregate effects are unclear	No
Soil nutrient pools	High	Fageria and Baligar (2005); Porter and Lawlor 1991 (p 173)	Nutrients are key limiting factors for yield (Boote <i>et al</i> 2013)	Yes
CO <sub>2</sub>	Low	Varotsos <i>et al</i> (2007)	Though CO <sub>2</sub> exerts a significant ecophysiological impact on crops (Long <i>et al</i> 2006, Leakey <i>et al</i> 2009, Sakurai <i>et al</i> 2014, Myers <i>et al</i> 2014), there is only low within-season variation	No
Soil salinity	Low	George <i>et al</i> (1997), Clarke <i>et al</i> (2002), Schofield and Kirkby (2003), Lambers <i>et al</i> (2008)	Could create variability in yield and production levels at spatially aggregated levels, but only low within-season variation (Ben-Ari and Makowski 2014)	No
Microorganisms	Unknown		May still be instrumental for understanding YV; management influences microorganisms (e.g. Gaudin <i>et al</i> 2015)	No
Soil acidity or density	Unknown		No information on its interannual variability is available	No



also has a possibly negative influence on soil micro-organism composition and on the severity of diseases (Hatfield *et al* 2011, Ahmed *et al* 2013).

The yield amount of wheat, maize and rice is reduced with non-optimal temperatures. Early growth, photosynthesis, carbon assimilation, stomatal conductance, plant development and root functioning strongly respond to temperature. This can diminish yields if temperature is too high or low (Schnyder 1993, Acevedo *et al* 2002, Wahid *et al* 2007, Barnabás *et al* 2008, Craufurd and Wheeler 2009, Farooq *et al* 2009a, 2011, Hatfield *et al* 2011, Hasanuzzaman *et al* 2013, Madhu and Hatfield 2013, Jagadish *et al* 2014, Suzuki *et al* 2014). Reproduction, again, is particularly sensitive to temperature extremes (Ishag and Mohamed 1996, Morison and Lawlor 1999, Dupont and Altenbach 2003, Barnabás *et al* 2008, Farooq *et al* 2011, Siebenmorgen *et al* 2013, Jagadish *et al* 2014). Many biochemical processes, like cell respiration and division, leaf senescence or membrane functionality depend on an optimal temperature range (Fuhrer 2003, Wahid *et al* 2007, Farooq *et al* 2009a, Mohammed and Tarpley 2009, Yadav 2010, Farooq *et al* 2011, Hasanuzzaman *et al* 2013, Miura and Furumoto 2013, Jagadish *et al* 2014). High T can also be coupled to an increased O<sub>3</sub> concentration that causes damage on its own (see below).

### 3.2.3. Processes affected by other important drivers

Solar radiation is the only source of energy for photosynthesis. Radiation is also the ultimate source of all weather variables like temperature. But the relation between radiation and temperature has recently become more complex (Wang and Dickinson 2013), and solar radiation affects crops in additional, distinct ways (Porter and Lawlor 1991 (p 106)). Excess radiation can damage the photosynthetic apparatus or induce oxidative stress, which both reduce the assimilation of C (Reddy *et al* 2004, Lambers *et al* 2008 (p 36)). Low radiation can also limit the uptake of nutrients (Lambers *et al* 2008 (p 268)).

Tropospheric ozone (O<sub>3</sub>) is known to cause substantial harm to crops in many regions (Avnery *et al* 2011, McGrath *et al* 2015). Its concentration in the Northern Hemisphere has risen in recent decades, with regional variation (Hoshika *et al* 2015). Increased [O<sub>3</sub>] has been shown to enhance leaf senescence, to impair reproductive processes and to lower the resistance against diseases (Fuhrer 2003, 2009, Hatfield *et al* 2011, Beckles and Thitisaksakul 2014). Higher [O<sub>3</sub>] can also counterbalance a fertilization effect of CO<sub>2</sub> (Fuhrer 2009, Hatfield *et al* 2011).

Nutrients including nitrogen (N), phosphorus (P) and other micronutrients are essential determinants of crop yield. Their uptake is influenced by temperature, soil characteristics (water content, acidity, salinity), root structure, soil characteristics, weed competition and plant growth (Fuhrer 2003, Barnabás *et al* 2008, Brouder and Volenec 2008, Ahmed *et al* 2013,

Ashraf 2014). Nutrients, especially N, and micronutrients like potassium or iron are required for photosynthesis, protein or starch synthesis, stress tolerance, turgor maintenance or ROS scavenging (Porter and Lawlor 1991 (p 13, 39, 55ff.); Hay and Porter 2006 (p 109, 198f); Thitisaksakul *et al* 2012, Powell *et al* 2012, Suzuki *et al* 2014). An excess of nutrients, in contrast, can cause misguided growth or impede grain filling (Schnyder 1993, Yang and Zhang 2006).

### 3.2.4. Influences on yield quality

Not only yield amount, but also yield quality is variable (e.g. Larcher 1995 (p 289); Dupont and Altenbach 2003, Siebenmorgen *et al* 2013). The assessment described above for yield amount has been performed for yield quality, too. It indicates that essentially the same set of drivers and plant physiological processes is important for the determination of quality (SI figure S6).


## 3.3. Selecting processes for improving crop models

The drivers of YV identified above affect yields indirectly by their influence on plant physiological processes. With our network we can also identify the relative importance of these processes to guide further development of crop models.

### 3.3.1. Relative importance of plant physiological processes

The putatively most important processes to transmit variability in growing conditions to YV are those strongly influenced by environmental or management stimuli and at the same time exerting a considerable impact on yields. An ordering diagram is shown in figure 5, which is structured as described in figure 2. Every process is located according to its sensitivity to driver variables on the *x*-axis and its respective impact on yield levels on the *y*-axis. We identify the following plant physiological processes as important (located in the top right quadrant): plant growth (split into growth of stem, roots and leaves); the uptake of water and nutrients; photosynthesis; reproduction; stress responses including antioxidant and disease defense; (evapo)transpiration; respiration; cell-internal processes like protein synthesis, turgor maintenance or division; senescence; and stomatal opening regulation. Connections between these processes, their environmental effectors and their influence on yields are found in section 3.2.

Three physiological hierarchy levels of the processes are included in the network: cell level, tissue and whole plant. Processes on the cell level determine those on tissue level (e.g. photosynthesis is required for root growth), and these in turn determine the processes on plant level (e.g. roots define the uptake of water). Growing conditions affect crops on different levels, which is thus reflected in our network.



There are minor differences between the three crops: wheat exhibits the full set of seventeen processes mentioned above as important, while maize and rice each have three less (senescence, stem development and stomatal opening regulation); SI figure S8. Different thresholds for the maximum allowed interaction path length do not alter the results qualitatively (SI figure S5).

### 3.3.2. Suggestions for implementing new features in crop models

To support the implementation of new features in crop models we collocate improvement suggestions derived from network structure and importance assessment. We compare the processes identified as important in our network analysis with the status quo of current crop models, as summarized by Boote *et al* (2013). Table 2 compiles this information for each plant physiological process (40 in total, on different hierarchical levels): importance for YV as ranked by the network evaluation (column 2), its implementation priority defined by Boote *et al* (2013) (col. 3), whether important drivers from the growing conditions *directly* influence the process (col. 4–9), and implementation suggestions (last column).

The processes identified as important by our network analysis mostly coincide with the priorities recommended by Boote *et al* (2013). Both sources rank

plant growth (in particular roots and leaves plus carbon allocation), reproductive processes including grain filling, the regulation of stomata and canopy energy balance, the nutrient balance, leaf senescence, respiration and photosynthesis (source-sink relationships) as priority for improving crop models. But differences in priority also occur in both directions. We identify cell turgor maintenance and protein synthesis as important, while these are not mentioned in Boote *et al* (2013). These two are usually not resolved in crop models, but rather covered by more coarse processes like water stress response or growth. Boote *et al* (2013), in contrast, rank grain filling, spikelet fertility and the response to pests and diseases as priorities for crop models. Yet we do not find these as primarily important processes in our network analysis. These processes are resolved in our network in the broader categories reproduction and stress responses, which are identified as important. The differences between our network method and the expert-approach by Boote *et al* (2013) are therefore mostly due to the network structure.

Interactions between different processes and the associated literature references for more details, like quantitative thresholds, can be easily extracted from the network. The full network is provided as GraphML source file in the SI for that purpose. Combined interactions between drivers or processes are particularly

**Table 2.** Plant physiological processes (first column) with their relative importance as our network suggests (c. 2), implementation priority according to our interpretation of Boote *et al* (2013) (c. 3; '1' is high, '2' is medium, and missing is unknown), environmental drivers for each process (only *direct* interactions; c. 4–9) according to our network, and options for improving current implementations (last c.; [B13] means they were also stated in Boote *et al* (2013)). Detailed interactions are listed in the SI. Abbreviations: Pr = Precipitation; SWC = Soil water content; T = Temperature; R<sub>s</sub> = Solar radiation; O<sub>3</sub> = Ozone; Nutr = Nutrient levels; SoilT = Soil temperature.

Process	Important (Network)	Priority (Boote <i>et al</i> 2013)	Drivers						Improvement options for crop models
			Pr/SWC	T	R <sub>s</sub>	[O <sub>3</sub> ]	Nutr	SoilT	
Plant dev. & growth	Yes	1	x	x	x	x		x	Inclusion of stressors in development timing: water deficit, T [B13] and salinity or O <sub>3</sub>
Roots growth	Yes	1	x	x				x	Growth response to edaphic [B13] and weather conditions
Leaves growth	Yes	2	x	x	x			x	Effects of canopy architecture, plant density and supply of nutrients and assimilates [B13]
Stem growth	Yes	2	x	x					Reserve accumulation and utilization under stress (T)
Early growth			x	x	x			x	Interactions of seed quality and stressors (water, T)
Reproduction	Yes	1	x	x	x	x			Fertility effects of high T, higher mechanistic detail [B13]
Grain filling		1	x	x		x	x		Stressors like high T [B13] and O <sub>3</sub> ; interactions with seed quality
Pollen development		1		x					Effects of high T [B13] and drought
Ovary development		1	x	x					Effects of heat and drought stress
Silking			x	x	x				Effects of heat and drought
Spikelet development			x	x					Effects of high T [B13] and water lack, interactions with elev. CO <sub>2</sub>
Anthesis		2	x	x					Effects of high T [B13] and radiation
Stomata regulation	Yes	1	x	x	x	x			Regulation by T, water, CO <sub>2</sub> [B13] and O <sub>3</sub> plus interaction with canopy effects and photosynthesis [B13]
Uptake	Yes								[See subprocess details]
Uptake of water		2	x	x					Interactions with root structure and nutrient uptake
Uptake of nutrients	Yes	1	x	x	x		x	x	Interactions with root structure [B13]; not only N but also P [B13]; effects of saline soils
General cell processes	Yes		x	x					[See subprocess details]
Division			x	x					Effects of drought and hot or cold T
Turgor maintenance	Yes		x	x					Drought and heat effects; interactions with growth and chemical reactions like photosynthesis

Table 2. (Continued.)

Process	Important (Network)	Priority (Boote <i>et al</i> 2013)	Drivers						Improvement options for crop models
			Pr/SWC	T	R <sub>s</sub>	[O <sub>3</sub> ]	Nutr	SoilT	
Protein synthesis	Yes			x	x				Impact on yield quality [B13]; effects of high T and radiation
Vernalization				x					Possible reversal under extreme heat
Senescence	Yes	1	x	x		x			Effects of high T [B13], drought, O <sub>3</sub> and excess fertilizer
Evapotranspiration	Yes	1		x	x				[See subprocess details]
Transpiration	Yes	1	x	x	x				Effects of elevated CO <sub>2</sub> and T, connection to photosynthesis [B13], effects of salinity
Evaporation				x	x				[no hints]
Respiration	Yes	2	x	x					Effects of CO <sub>2</sub> [B13] and extreme T
Photosynthesis	Yes	1	x	x	x	x			Scaling up from leaf to canopy or field [B13]; effects of non-optimal T or drought
Light interception									[no hints]
Production of ROS			x	x	x	x			Effects of extreme T and drought, and the impact of ROS on other processes like photosynthesis
Photorespiration			x	x					Effects of heat stress
Acclimation									[no hints]
Stress responses	Yes	2							[See subprocess details]
Antioxidant defense			x	x					Induction by high T or drought through increased ROS production
Sec. metabol. accum.				x					Energy costs by high T
Comp. osmol. accum.			x	x					Energy costs by drought or high T
Expr. of stress prot.				x					Energy costs by high T
Responses to diseases		2		x		x			Incorporation of diseases into crop models [B13]; interaction with energy balance
Escape or avoidance			x						[no hints]
Allelopathy									[no hints]
Herbivory defenses		2							Incorporation into crop models [B13] together with pest models; interaction with energy balance

relevant since plant responses to simultaneous changes in growing conditions often differ from the responses to individual changes (Lobell and Burke 2010, Jagadish *et al* 2014, Ray *et al* 2015). Since extreme events can induce nonlinear responses in crops, their impact on plant physiological processes is of particular importance. These influences are annotated explicitly for the network interactions where mentioned in the associated studies (full network in SI).

## 4. Discussion

### 4.1. Validity of the network method

A network structure, derived from literature, is employed to evaluate the importance of growing condition factors and plant physiological processes on YV. Hence it is eminent to have an unbiased knowledge base for its construction. With the systematic approach by pre-defined search terms we aim to keep a literature bias (i.e. the over-representation of aspects like temperature) at a minimum. In addition, all interaction edges have the same weight independent of how often they are confirmed (or contradicted) in the literature, which limits a potential research frequency bias. A strong representation of a process in the literature might, however, reflect its pertinency for implementation. Additionally, a broad literature coverage of aspects like heat or drought stress might stem from its agronomic importance—which further warrants their appropriate consideration in models. Therefore we argue that our findings, which are based on a large interaction network and are robust under different analysis setups, are relevant for crop models.

Our importance assessment does not consider quantitative information in the interactions. But for the relative weighting of process importance a quantitative network would not necessarily be more accurate, as it would introduce more parameters to the method. Furthermore, every quantitative parameter would depend on crop, cultivar and location—which would be beyond the scope of any single meta-study to curtail for 350 interactions. Breeding efforts have achieved higher sensitivities to selected growing conditions, e.g. N and water provision. This trend is neglected in our network, for the same reason of quantitative complexity, but we argue as above that the method would not necessarily benefit from its consideration. Another possible issue that comes with missing quantitative information is that a node with many small influences on other nodes is considered more important than a node with only few but large impacts. Yet many small impacts can also amount to large ones, and the quantity argument goes as above. But if necessary, quantitative information for any specific process can easily be retrieved from the recorded interaction references.

The network unites plant physiological processes on cell, tissue and whole plant level. Single, scale-dependent networks for each of these three would be an alternative approach that respects differences between levels. But we argue that a united approach is justified in our case for three reasons. First, small-scale processes (e.g. cell respiration) accumulate influence over the growing season and therefore can determine yields as much as large-scale processes like, for example, herbivory. Second, the network is constructed to deduce improvement suggestions for crop models, which also need to reflect yield influences on all three levels. Thus we can more easily derive these suggestions with a combined network, but endorse further differentiation in later work. Third, we explicitly aim to capture all relevant mechanisms that may or may not act synchronously to influence yields (synchrony as, for example, in vernalization).

Plant physiological processes can be grouped or aggregated in manifold ways (e.g. Boote *et al* 2013, Bassu *et al* 2014). Our network is therefore only one approach to classify these processes. It is elicited from sequential literature reading of plant physiology text books and independent articles. An assessment of how different basic network structures (differing from figure 3) would affect the results was not tested here. But the network proved flexible enough to incorporate all interactions and elements found in the literature. Some plant responses to growing conditions may not be included in the 350 interactions of our network. Yet we argue that these are likely only minor given the systematic literature mining and the robust driver ranking.

The analysis indicates temperature and precipitation as strong drivers of YV—which is well-known and thus confirms the validity of our method. But we have also identified further factors whose own variability could imply variability in yields. Drivers which our network approach labels as ‘unimportant’ are not necessarily unimportant in reality—the relative weighting applied here only assigns more weight to the others. In contrast, the drivers defined as important by the network structure have only the potential to cause variability. Yet the actual importance depends on the specific combination of the individual components of the growing conditions. Although CO<sub>2</sub> or soil salinity are not regarded as important contributors to YV, they can strongly influence responses to other stressors via interacting effects (Jagadish *et al* 2014). Adequately representing yield quality is equally judged an essential target for crop model improvement, evidenced by the 13 direct influences on yield quality in the network.

The close similarity of results for wheat, maize and rice arises from two independent factors. First, the network edges are often (68%) based on publications that are valid for all three crops considered. Second, the network is qualitative only such that quantitative distinctions between crops are not accessible. Differences between crops (e.g. rice is usually irrigated, maize is a C<sub>4</sub>



crop, winter wheat requires vernalization) are not questioned. But with regard to modeling the generality of the drivers identified is beneficial, since most of the mechanistic pathways are shared between crops.

The network essentially reflects the complexity of plant regulatory systems. This entails a high ‘importance’ for drivers or processes that are involved in several regulatory pathways, i.e. importance reflects complexity. Complex systems may either be prone to abrupt state changes under disturbances (Robbirt *et al* 2014, Willmer 2014, Zscheischler *et al* 2014, Franklin *et al* 2016) or enhance the stability of a system (an example is resilience from biodiversity). The current network does not reveal whether a process that influences YV actually enhances or dampens it. This requires deeper inspection of each single interaction with quantitative information. Our method suggests exactly these crucial points that need further inspection. One possible inspection approach is a recursive refinement of the network into single-process subsets.

#### 4.2. Applicability of the implementation suggestions

Although the network has been constructed without input from crop models we apply it to guide model improvement strategies. This approach is uncommon, but we adopted it for maintaining an ‘outside’ look on the models inspired by experimental literature alone. In addition the very diverse types and characteristics of models (Rosenzweig *et al* 2014, Elliott *et al* 2015) require an abstract approach that does not depend on a certain class of models. The agreement about the major improvement points between our network method and the expert approach by Boote *et al* (2013) show the efficacy of the network method to detect essential features from the literature. It also justifies the assumption that ‘importance’ can be derived from the number of connections in the network. Differences in priorities reflect the potential for supplementing one approach with the other.

Many of the process improvement options for crop models are targeting currently less well represented physiological interactions. These general suggestions have to be adjusted for each particular model. There is no guarantee that a crop model eventually becomes better in modeling yield (variability) with a finer resolution of processes or by adding new ones. More processes require more parameters, which could entail model or calibration errors. The necessary experimental data are not easy to find, but one possible starting point is the AgTrials database<sup>4</sup>. Some processes may not yet be implemented for their high complexity paired with unclear benefits for the model. Examples are the crop responses to pests and diseases (high specificity of crop–pathogen–environment interaction; Luck *et al* 2011) or to an elevated ozone concentration (lack of global databases, unclear effect on aggregate level). Nonetheless these have potential

to help understanding of YV in diverse environments, and from sources other than temperature or precipitation. Regional-scale crop patterns have been studied as causes of YV by Ben-Ari and Makowski (2014), and genetic traits in YV by Mickelbart *et al* (2015). The focus on plant physiological process level in our analysis complements these two approaches.

## 5. Conclusion

We have applied a novel methodology for a systematic literature review to identify and rank the importance of drivers and plant physiological processes for crop YV. We have also derived a comprehensive list of target points for improving crop models with respect to YV. As expected, our method confirms that current modeling approaches have addressed many of the important drivers of YV. Thus our approach can be seen as a cross-validation of existing modeling concepts. However, we also show that the drivers and the mechanisms implemented are not sufficiently comprehensive, which thus can guide future model improvement. Our network is a unique structured summary of the literature and its free accessibility can support the improved representation of YV in crop models. In particular the network interaction structure and the rich quantitative literature information associated with it can serve as a starting point.

The approach could be extended by a semi-automatic text mining, extracting the most relevant information from literature databases. Text mining has successfully been applied in medical bioinformatics (Zhu *et al* 2013, Fluck and Hofmann-Apitius 2014, Fleuren and Alkema 2015). Our network-based review could serve as a first step towards this. We have shown its methodical validity as a structured, consistent and scalable dictionary of literature knowledge. The approach is easily applicable to many other fields of research.

## Acknowledgments

The authors want to thank Frank Wechsung, Christopher Reyer, Christoph Gornott, Katharina Waha (for helpful discussions) and Deepak Ray (for data provision). BS acknowledges financial support from the German National Academic Foundation, SR from the MACSUR project (031A103B) funded by the German Federal Ministry of Education and Research (BMBF), and CM from the MACMIT project (01LN1317A) funded by BMBF.

## References

- Acevedo E, Silva P and Silva H 2002 Wheat growth and physiology ([www.fao.org/docrep/006/y4011e/y4011e06.htm#bm06](http://www.fao.org/docrep/006/y4011e/y4011e06.htm#bm06)) (Accessed: 21 November 2016)
- Ahmed F, Rafii M Y, Ismail M R, Juraimi A S, Rahim H A, Asfaliza R and Latif M A 2013 Waterlogging tolerance of crops: breeding, mechanism of tolerance, molecular

<sup>4</sup> <http://agtrials.org/>; accession date 01 February 2016.



- approaches, and future prospects *BioMed Res. Int.* **2013** 963525
- Ashraf M 2014 Stress-induced changes in wheat grain composition and quality *Crit. Rev. Food Sci. Nutrition* **54** 1576–83
- Asseng S et al 2015 Rising temperatures reduce global wheat production *Nat. Clim. Change* **5** 143–7
- Asseng S, Foster I and Turner N C 2011 The impact of temperature variability on wheat yields *Glob. Change Biol.* **17** 997–1012
- Avnery S, Mauzerall D L, Liu J and Horowitz L W 2011 Global crop yield reductions due to surface ozone exposure: I. Year 2000 crop production losses and economic damage *Atmos. Environ.* **45** 2284–96
- Bakker M M, Govers G, Ewert F, Rounsevell M and Jones R 2005 Variability in regional wheat yields as a function of climate, soil and economic variables: Assessing the risk of confounding *Agric. Ecosyst. Environ.* **110** 195–209
- Barlow K, Christy B, O'Leary G, Riffkin P and Nuttall J 2015 Simulating the impact of extreme heat and frost events on wheat crop production: a review *Field Crops Res.* **171** 109–19
- Barnabás B, Jäger K and Fehér A 2008 The effect of drought and heat stress on reproductive processes in cereals *Plant Cell Environ.* **31** 11–38
- Bassu S et al 2014 How do various maize crop models vary in their responses to climate change factors? *Glob. Change Biol.* **20** 2301–20
- Beckles D M and Thitisaksakul M 2014 How environmental stress affects starch composition and functionality in cereal endosperm *Starch* **66** 58–71
- Bell J E, Sherry R and Luo Y 2010 Changes in soil water dynamics due to variation in precipitation and temperature: an ecohydrological analysis in a tallgrass prairie *Water Resour. Res.* **46** 1–11
- Ben-Ari T and Makowski D 2014 Decomposing global crop yield variability *Environ. Res. Lett.* **9** 114011
- Blois J L, Williams J W, Fitzpatrick M C, Jackson S T and Ferrier S 2013 Space can substitute for time in predicting climate-change effects on biodiversity *Proc. Natl Acad. Sci.* **110** 9374–9
- Boote K J, Jones J W, White J W, Asseng S and Lizaso J I 2013 Putting mechanisms into crop production models *Plant Cell Environ.* **36** 1658–72
- Boyer J S and Westgate M E 2004 Grain yields with limited water *J. Exp. Bot.* **55** 2385–94
- Brouder S M and Volenec J J 2008 Impact of climate change on crop nutrient and water use efficiencies *Physiol. Plantarum* **133** 705–24
- Chung U, Gbegbelegbe S, Shiferaw B, Robertson R, Yun J J, Tesfaye K, Hoogenboom G and Sonder K 2014 Modeling the effect of a heat wave on maize production in the USA and its implications on food security in the developing world *Weather Clim. Extremes* **5–6** 67–77
- Clarke C J, George R J, Bell R W and Hatton T J 2002 Dryland salinity in south-western Australia: its origins, remedies, and future research directions *Aust. J. Soil Res.* **40** 93–113
- Coumou D and Rahmstorf S 2012 A decade of weather extremes *Nat. Clim. Change* **2** 1–6
- Craufurd P Q and Wheeler T R 2009 Climate change and the flowering time of annual crops *J. Exp. Bot.* **60** 2529–39
- de Bossoreille de Ribou S, Douam F, Hamant O, Frohlich M W and Negruțiu I 2013 Plant science and agricultural productivity: why are we hitting the yield ceiling? *Plant Sci.* **210** 159–76
- Deryng D, Conway D, Ramankutty N, Price J and Warren R 2014 Global crop yield response to extreme heat stress under multiple climate change futures *Environ. Res. Lett.* **9** 034011
- Donat M G et al 2013 Updated analyses of temperature and precipitation extreme indices since the beginning of the twentieth century: the HadEX2 dataset *J. Geophys. Res.: Atmos.* **118** 2098–118
- Dupont F and Altenbach S 2003 Molecular and biochemical impacts of environmental factors on wheat grain development and protein synthesis *J. Cereal Sci.* **38** 133–46
- Elliott J et al 2015 The Global Gridded Crop Model intercomparison: data and modeling protocols for Phase 1 (v1.0) *Geosci. Model Dev.* **7** 4383–427
- Eyshi Rezaei E, Webber H, Gaiser T, Naab J and Ewert F 2014 Heat stress in cereals: mechanisms and modelling *Eur. J. Agronomy* **64** 98–113
- Fageria N K and Baligar V C 2005 Enhancing nitrogen use efficiency in crop plants *Adv. Agronomy* **88** 97–185
- FAO 2015 <http://faostat3.fao.org/home/E> (Accessed 21 July 2015)
- Farooq M, Aziz T, Wahid A, Lee D-j and Siddique K H M 2009a Chilling tolerance in maize: agronomic and physiological approaches *Crop Pasture Sci.* **60** 501–16
- Farooq M, Bramley H, Palta J A and Siddique K H M 2011 Heat stress in wheat during reproductive and grain-filling phases *Crit. Rev. Plant Sci.* **30** 491–507
- Farooq M, Hussain M and Siddique K H M 2014 Drought stress in wheat during flowering and grain-filling periods *Crit. Rev. Plant Sci.* **33** 331–49
- Farooq M, Wahid A, Kobayashi N, Fujita D and Basra S M A 2009b Plant drought stress: effects, mechanisms and management *Agronomy Sustain. Dev.* **29** 185–212
- Fleuren W W and Alkema W 2015 Application of text mining in the biomedical domain *Methods* **74** 97–106
- Fluck J and Hofmann-Apitius M 2014 Text mining for systems biology *Drug Discovery Today* **19** 140–4
- Franklin J, Serra-diaz J M, Syphard A D, Regan H M and Framework C 2016 Global change and terrestrial plant community dynamics *Proc. Natl Acad. Sci. USA* **113** 3725–34
- Fuhrer J 2003 Agroecosystem responses to combinations of elevated CO<sub>2</sub>, ozone, and global climate change *Agric. Ecosyst. Environ.* **97** 1–20
- Fuhrer J 2009 Ozone risk for crops and pastures in present and future climates *Naturwissenschaften* **96** 173–94
- Gaudin A C M, Tolhurst T N, Ker A P, Janovicek K, Tortora C, Martin R C and Deen W 2015 Increasing crop diversity mitigates weather variations and improves yield stability *Plos One* **10** e0113261
- George R, McFarlane D and Nulsen B 1997 Salinity threatens the viability of agriculture and ecosystems in Western Australia *Hydrogeol. J.* **5** 6–21
- Gonzalez-Dugo V, Durand J-L and Gastal F 2010 Water deficit and nitrogen nutrition of crops a review *Agronomy Sustain. Dev.* **30** 529–44
- Hall D O, Scurlock J M O, Bolhar-Nordenkamp H R, Leegood R C and Long S P (ed) 1993 *Photosynthesis and Production in a Changing Environment* 1st edn (London: Chapman and Hall)
- Hasanuzzaman M, Nahar K, Alam M M, Roychowdhury R and Fujita M 2013 Physiological, biochemical, and molecular mechanisms of heat stress tolerance in plants *Int. J. Mol. Sci.* **14** 9643–84
- Hatfield J L, Boote K J, Kimball B A, Ziska L H, Izaurralde R C, Ort D, Thomson A M and Wolfe D 2011 Climate impacts on agriculture: implications for crop production *Agronomy J.* **103** 351–70
- Hay R K M and Porter J R 2006 *The Physiology of Crop Yield* 2nd edn (Oxford: Blackwell)
- Hay R K M and Walker A J 1989 *An Introduction to the Physiology of Crop Yield* 1st edn (Essex: Longman Scientific & Technical)
- Headey D and Fan S 2008 Anatomy of a crisis: the causes and consequences of surging food prices *Agric. Econ.* **39** 375–91
- Headey D D 2010 Rethinking the global food crisis: the role of trade shocks *IFPRI institute discussion paper 00958* (<https://core.ac.uk/download/pdf/6250134.pdf>)
- Holzworth D P, Snow V, Janssen S, Athanasiadis I N, Donatelli M, Hoogenboom G, White J W and Thorburn P 2015 Agricultural production systems modelling and software: current status and future prospects *Environ. Model. Softw.* **2014** 1–11
- Hoshika Y, Katata G, Deushi M, Watanabe M, Koike T and Paoletti E 2015 Ozone-induced stomatal sluggishness changes carbon and water balance of temperate deciduous forests *Nat. Sci. Rep.* **5** 9871
- Ishag H M and Mohamed B A 1996 Phasic development of spring wheat and stability of yield and its components in hot environments *Field Crops Res.* **46** 169–76

- Jagadish K S V, Niteen N K, Gui X, Reneeliza J M, Rajeev N B, Cheryl Q, Anandhan T and Prasad P V V 2014 Agronomic and physiological responses to high temperature, drought, and elevated CO<sub>2</sub> interactions in cereals *Advances in Agronomy* ed D L Sparks 127 edn (Amsterdam: Elsevier) pp 111–56
- Lambers H, Chapin F S and Pons T L 2008 *Plant Physiological Ecology* 2nd edn (New York: Springer)
- Larcher W 1995 *Physiological Plant Ecology* 3rd edn (New York: Springer)
- Lawlor D W and Tezara W 2009 Causes of decreased photosynthetic rate and metabolic capacity in water-deficient leaf cells: a critical evaluation of mechanisms and integration of processes. *Ann. Bot.* **103** 561–79
- Leakey A D B, Ainsworth E A, Bernacchi C J, Rogers A, Long S P and Ort D R 2009 Elevated CO<sub>2</sub> effects on plant carbon, nitrogen, and water relations: six important lessons from FACE *J. Exp. Bot.* **60** 2859–76
- Lesk C, Rowhani P and Ramankutty N 2016 Influence of extreme weather disasters on global crop production *Nature* **529** 84–7
- Lobell D B and Burke M B 2010 On the use of statistical models to predict crop yield responses to climate change *Agric. Forest Meteorol.* **150** 1443–52
- Lobell D B and Gourdji S M 2012 The influence of climate change on global crop productivity *Plant Physiol.* **160** 1686–97
- Lobell D B, Hammer G L, McLean G, Messina C, Roberts M J and Schlenker W 2013 The critical role of extreme heat for maize production in the United States *Nat. Clim. Change* **3** 497–501
- Long S P, Ainsworth E A, Leakey A D B, Nösberger J and Ort D R 2006 Food for thought: lower-than-expected crop yield stimulation with rising CO<sub>2</sub> concentrations *Science* **312** 1918–21
- Luck J, Spackman M, Freeman A, Trebicki P, Griffiths W, Finlay K and Chakraborty S 2011 Climate change and diseases of food crops *Plant Pathol.* **60** 113–21
- Madhu M and Hatfield J L 2013 Dynamics of plant root growth under increased atmospheric carbon dioxide *Agronomy J.* **105** 657
- Martiello M A and Giacchi M V 2010 High temperatures and health outcomes: a review of the literature *Scand. J. Public Health* **38** 826–37
- Martre P et al 2015 Multimodel ensembles of wheat growth: many models are better than one *Glob. Change Biol.* **21** 911–25
- McGrath J M, Betzelberger A M, Wang S, Shook E, Zhu X-G, Long S P and Ainsworth E A 2015 An analysis of ozone damage to historical maize and soybean yields in the United States *Proc. Natl Acad. Sci.* **112** 14390–5
- Mickelbart M V, Hasegawa P M and Bailey-Serres J 2015 Genetic mechanisms of abiotic stress tolerance that translate to crop yield stability *Nat. Rev. Genet.* **16** 237–51
- Miura K and Furumoto T 2013 Cold signaling and cold response in plants *Int. J. Mol. Sci.* **14** 5312–37
- Mohammed A-R and Tarpley L 2009 Impact of High Nighttime Temperature on Respiration, Membrane Stability, Antioxidant Capacity, and Yield of Rice Plants *Crop Sci.* **49** 313
- Morison J I L and Lawlor D W 1999 Interactions between increasing CO<sub>2</sub> concentration and temperature on plant growth *Plant Cell Environ.* **22** 659–82
- Morton J F 2007 The impact of climate change on smallholder and subsistence agriculture *Proc. Natl Acad. Sci. USA* **104** 19680–5
- Myers S S et al 2014 Increasing CO<sub>2</sub> threatens human nutrition *Nature* **510** 139–42
- Palosuo T et al 2011 Simulation of winter wheat yield and its variability in different climates of Europe: a comparison of eight crop growth models *Eur. J. Agronomy* **35** 103–14
- Porter J R and Lawlor D W (ed) 1991 *Plant Growth Interactions with Nutrition and Environment* 1st edn (Cambridge: Cambridge University Press)
- Porter J R and Semenov M A 2005 Crop responses to climatic variation *Phil. Trans. R. Soc. B* **360** 2021–35
- Portmann F T, Siebert S and Döll P 2010 MIRCA2000-global monthly irrigated and rainfed crop areas around the year 2000: a new high-resolution data set for agricultural and hydrological modeling *Glob. Biogeochem. Cycles* **24** 1–24
- Powell N, Ji X, Ravash R, Edlington J and Dolferus R 2012 Yield stability for cereals in a changing climate *Funct. Plant Biol.* **39** 539–52
- Rahmstorf S and Coumou D 2011 Increase of extreme events in a warming world *Proc. Natl Acad. Sci.* **108** 17905–9
- Ray D K, Gerber J S, MacDonald G K and West P C 2015 Climate variation explains a third of global crop yield variability *Nat. Commun.* **6** 5989
- Reddy A R, Chaitanya K V and Vivekanandan M 2004 Drought-induced responses of photosynthesis and antioxidant metabolism in higher plants *J. Plant Physiol.* **161** 1189–202
- Robbirt K M, Roberts D L, Hutchings M J and Davy A J 2014 Potential disruption of pollination in a sexually deceptive orchid by climatic change *Curr. Biol.* **24** 2845–9
- Rosenzweig C et al 2014 Assessing agricultural risks of climate change in the 21st century in a global gridded crop model intercomparison *Proc. Natl Acad. Sci.* **111** 3268–73
- Rötter R P 2014 Robust uncertainty *Nat. Clim. Change* **4** 251–2
- Rötter R P, Carter T R, Olesen J E and Porter J R 2011 Crop-climate models need an overhaul *Nat. Clim. Change* **1** 175–7
- Ruane A C, Goldberg R and Chrysanthacopoulos J 2015 Climate forcing datasets for agricultural modeling: merged products for gap-filling and historical climate series estimation *Agric. Forest Meteorol.* **200** 233–48
- Sakurai G, Iizumi T, Nishimori M and Yokozawa M 2014 How much has the increase in atmospheric CO<sub>2</sub> directly affected past soybean production? *Sci. Rep.* **4** 1–5
- Sánchez B, Rasmussen A and Porter J R 2014 Temperatures and the growth and development of maize and rice: a review *Glob. Change Biol.* **20** 408–17
- Schlenker W and Roberts M J 2009 Nonlinear temperature effects indicate severe damages to U.S. crop yields under climate change *Proc. Natl Acad. Sci. USA* **106** 15594–8
- Schmidhuber J and Tubiello F N 2007 Global food security under climate change *Proc. Natl Acad. Sci. USA* **104** 19703–8
- Schnyder H 1993 The role of carbohydrate storage and redistribution in the source-sink relations of wheat and barley during grain filling—a review *New Phytologist* **123** 233–45
- Schofield R V and Kirkby M J 2003 Application of salinization indicators and initial development of potential global soil salinization scenario under climatic change *Glob. Biogeochem. Cycles* **17** 1–13
- Seneviratne S et al 2012 Changes in climate extremes and their impacts on the natural physical environment *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation: A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change (IPCC)* ed C Field et al (Cambridge: Cambridge University Press) pp 109–230
- Siebenmorgen T J, Grigg B C and Lanning S B 2013 Impacts of preharvest factors during kernel development on rice quality and functionality *Annu. Rev. Food Sci. Technol.* **4** 101–15
- Suzuki N, Rivero R M, Shulaev V, Blumwald E and Mittler R 2014 Abiotic and biotic stress combinations *New Phytologist* **203** 32–43
- Tai A P K, Martin M V and Heald C L 2014 Threat to future global food security from climate change and ozone air pollution *Nat. Clim. Change* **4** 817–21
- Thitisaksakul M, Jiménez R C, Arias M C and Beckles D M 2012 Effects of environmental factors on cereal starch biosynthesis and composition *J. Cereal Sci.* **56** 67–80
- Thornton P K, Ericksen P J, Herrero M and Challinor A J 2014 Climate variability and vulnerability to climate change: a review *Glob. Change Biol.* **20** 3313–28
- Tokatlidis I S 2014 Addressing the yield by density interaction is a prerequisite to bridge the yield gap of rain-fed wheat *Ann. Appl. Biol.* **165** 27–42
- Tollenaar M and Lee E A 2002 Yield potential, yield stability and stress tolerance in maize *Field Crops Res.* **75** 161–9
- Varotsos C, Assimakopoulos M N and Efsthathiou M 2007 Technical note: long-term memory effect in the atmospheric CO<sub>2</sub>

- concentration at Mauna Loa *Atmos. Chem. Phys. Discuss.* **6** 11957–70
- Waha K, Müller C, Bondeau A, Dietrich J, Kurukulasuriya P, Heinke J and Lotze-Campen H 2013 Adaptation to climate change through the choice of cropping system and sowing date in sub-Saharan Africa *Glob. Environ. Change* **23** 130–43
- Waha K, van Bussel L G J, Müller C and Bondeau A 2012 Climate-driven simulation of global crop sowing dates *Glob. Ecol. Biogeogr.* **21** 247–59
- Wahid A, Gelani S, Ashraf M and Foolad M 2007 Heat tolerance in plants: an overview *Environ. Exp. Bot.* **61** 199–223
- Wang K and Dickinson R E 2013 Contribution of solar radiation to decadal temperature variability over land *Proc. Natl Acad. Sci. USA* **110** 14877–82
- Wheeler T and von Braun J 2013 Climate change impacts on global food security *Science* **341** 508–13
- White J W, Hoogenboom G, Kimball B A and Wall G W 2011 Methodologies for simulating impacts of climate change on crop production *Field Crops Res.* **124** 357–68
- Wild O et al 2012 Modelling future changes in surface ozone: a parameterized approach *Atmos. Chem. Phys.* **12** 2037–54
- Willmer P 2014 Climate change: bees and orchids lose touch *Curr. Biol.* **24** R1133–5
- Yadav S K 2010 Cold stress tolerance mechanisms in plants: a review *Agronomy Sustain. Dev.* **30** 515–27
- Yang J and Zhang J 2006 Grain filling of cereals under soil drying *New Phytologist* **169** 223–36
- Yu Q, Li L, Luo Q, Eamus D, Xu S, Chen C, Wang E, Liu J and Nielsen D C 2014 Year patterns of climate impact on wheat yields *Int. J. Climatol.* **34** 518–28
- Zhu F, Patumcharoenpol P, Zhang C, Yang Y, Chan J, Meechai A, Vongsangnak W and Shen B 2013 Biomedical text mining and its applications in cancer research *J. Biomed. Inform.* **46** 200–11
- Zscheischler J et al 2014 A few extreme events dominate global interannual variability in gross primary production *Environ. Res. Lett.* **9** 035001